arXiv:1503.03927v1 [math.DS] 13 Mar 2015

Multiple Forced rotations
for the N-pendulum Equation

Hui Qiao

School of Mathematics and Statistics

Wuhai University, Wuhan 430072,

The People's Republic of China

e-mail: qiaohuimath@whu.edu.cn

Abstract

In this paper, we consider the planar forced N-pendulum equation. Multiple rotational solutions are obtained.

2010 Mathematics Subject Classification. 37J45, 34C25, 58E05.

Key words. critical points, Lagrangian systems, rotations, multiplicity, composed pendulum.

1 Introduction and main results

Let m_i, ℓ_i be positive constants, $i = 1,2,\cdots,N$, and

$$\alpha_j = \sum_{s=j}^{N} m_s,\ \beta_j = \alpha_j \ell_j,\ j = 1,2,\cdots,N. \tag{1.1}$$

The motion of the coplanar N-pendulum is governed by the Lagrangian

$$L(q,p) = \frac{1}{2}A(q)p\cdot p + V(q), \tag{1.2}$$

where

$$q = (q_1,q_2,\cdots,q_N),\ p = (p_1,p_2,\cdots,p_N) \in \mathbb{R}^N,$$

$$A(q)p\cdot p = \sum_{1\leqslant i\leqslant N} \alpha_i \ell_i^2 p_i^2 + \sum_{1\leqslant i<j\leqslant N} 2\alpha_j \ell_i \ell_j \cos(q_i - q_j) p_i p_j, \tag{1.3}$$

$$V(q) = g\sum_{j=1}^{N} \beta_j \cos(q_j), \tag{1.4}$$

In the following, we add an periodic forcing term $f(t) = (f_1(t), \cdots, f_N(t))$. The corresponding system of equations are

$$\frac{\mathrm{d}}{\mathrm{d}t} L_p(q,\dot{q}) - L_q(q,\dot{q}) = f(t), \tag{1.5}$$

Periodic motions for the N-pendulum are solutions of (1.5) satisfying

$$q(t+T) - q(t) = 2\pi v,\ v \in \mathbb{Z}^N,\ \forall t \in \mathbb{R}. \tag{1.6}$$

In (1.6), if $v = 0$, solutions are oscillatory. Existences and multiplicities of oscillations for planar double, triple and N-pendulum have been studied in [3, 8, 4, 17, 15] and references therein. In [14], Rabinowitz showed the existence of at least $N+1$ 2π-periodic forced oscillations for more general Lagrangian functions.

In this paper, we consider solutions with

$$v = (v_1, v_2, \cdots, v_N) \neq 0,$$

which are called *rotational solutions* and non-contractible on T^N. Note that if $q(t)$ is a solution, then $q(kT+t) - q(t) \equiv kv$. We restrict choices of rotational vectors to $\mathbb{Z}_1^N$ defined as follows.

Definition 1. A rotational vector $v = (v_1, v_2, \cdots, v_N) \in \mathbb{Z}^N \setminus \{0\}$ is called *prime*, if one of its coordinates equals to 1 while others are zero, or two of them are relatively prime. Denotes by $\mathbb{Z}_1^N$ the set of all prime rotational vectors in $\mathbb{Z}^N$.

For $v \in \mathbb{Z}_1^N$, let

$$q(t) = x(t) + \frac{2\pi v t}{T},\ \dot{q}(t) = \dot{x}(t) + \frac{2\pi v}{T}, \tag{1.7}$$

and define a functional $\mathscr{L} : E = W^{1,2}(S^1, \mathbb{R}^N) \to \mathbb{R}$ by

$$\mathscr{L}(x) = \int_0^T \big(L(q(t), \dot{q}(t)) + f(t) \cdot q(t)\big)\mathrm{d}t. \tag{1.8}$$

Its critical points give rotational solutions. When we study multiplicities, the following symmetries are involved:

$\mathbb{Z}^N$-*symmetry*. Note that E can be orthogonally decomposed as

$$E = E^0 \oplus \widetilde{E}, \tag{1.9}$$

where E^0 denotes the subspace of constant functions and $\widetilde{E}$ denotes the subspace of functions with zero mean value. Since the function L is 2π-periodic in q_i for all $1 \leqslant i \leqslant N$ and $f \in \widetilde{E}$, the functional $\mathscr{L}$ defined by (1.8) is also 2π-periodic in all q_is. Hence it can be defined on

$$\mathscr{E} = E/(2\pi\mathbb{Z}^N) = T^N \times \widetilde{E}. \tag{1.10}$$

S^1-*symmetry*. If $f \equiv 0$, there is an S^1-action involved in the problem. Given $v \in \mathbb{Z}_1^N$ and $T > 0$, let

$$\theta \cdot x(t) = x(t+\theta) + \frac{2\pi v}{T}\theta, \tag{1.11}$$

where we use the representation of S^1 by $\left\{\exp\left(\frac{2\pi\sqrt{-1}}{T}\theta\right) \;\middle|\; \theta \in [0,T]\right\}$. Then $\mathscr{L}$ is S^1-invariant. Moreover, this action is free.

Denote by $\mathscr{P}(v,T)$ the set of distinct rotational solutions of (1.5) with given rotational vector v and period T.

Theorem 1. *Assume that f satisfies*

$$f \in L^2(\mathbb{R}/(T\mathbb{Z}),\mathbb{R}^N), \ \int_0^T f(t)dt = 0. \tag{1.12}$$

For every $v \in \mathbb{Z}_1^N$ and $T > 0$, we have
(i) If $f \not\equiv 0$, then

$$^{\#}\mathscr{P}(v,T) \geqslant N+1.$$

If all solutions are nondegenerate, then $^{\#}\mathscr{P}(v,T) \geqslant 2^N$.
(ii) If $f \equiv 0$, then

$$^{\#}\mathscr{P}(v,T) \geqslant N.$$

If all solutions are nondegenerate, then $^{\#}\mathscr{P}(v,T) \geqslant 2^{N-1}$.

Similarly to [8, 4], the aim of this paper is to establish additional rotational solutions for special arrangement of masses m_j and lengths ℓ_j instead of the nondegenerate assumption. In the following result, we consider $v = (v_1, v_2, \cdots, v_N) \in \mathbb{Z}_1^N$ with zero components. Let

$$N_0 = {}^{\#}\{i \mid 1 \leqslant i \leqslant N, \ v_i = 0\}. \tag{1.13}$$

Theorem 2. *Assume that f satisfies* (1.12) *and*

$$|f(t)| \leqslant M_0, \ \forall t \in \mathbb{R}. \tag{1.14}$$

If $1 \leqslant N_0 \leqslant N-1$, there exist masses m_j, lengths ℓ_j ($1 \leqslant j \leqslant N$), and constants $T_1, T_2 > 0$ such that for each $T \in [T_1, T_2]$, we have
(i) If $f \not\equiv 0$, then

$$^{\#}\mathscr{P}(v,T) \geqslant (N - N_0 + 1)2^{N_0}.$$

(ii) If $f \equiv 0$, then

$$^{\#}\mathscr{P}(v,T) \geqslant (N - N_0)2^{N_0}.$$

2 Variational settings

For $x = (x_1, x_2, \cdots, x_N) \in \mathscr{E}$, denote by $||x||$ the standard L^2-norm of $x(t)$, i.e.,

$$||x||^2 = \int_0^T |x(t)|^2 \mathrm{d}t = \sum_{i=1}^N \int_0^T x_i^2(t) \mathrm{d}t.$$

The functional $\mathscr{L} : \mathscr{E} \to \mathbb{R}$ defined by (1.8) can be written as

$$\mathscr{L}(x) = \mathscr{L}_1(x) + \mathscr{L}_2(x) + \mathscr{L}_3(x), \tag{2.1}$$

where

$$\mathscr{L}_1(x) = \frac{1}{2}\langle A(q)\dot{q}, \dot{q}\rangle,\ \mathscr{L}_2(x) = \int_0^T V(q)\mathrm{d}t,\ \mathscr{L}_3(x) = \int_0^T f(t)\cdot q\mathrm{d}t. \tag{2.2}$$

According to (1.10), we write $x \in \mathscr{E}$ as

$$x = \bar{x} + \widetilde{x}, \quad \bar{x} = (\bar{x}_1, \bar{x}_2, \cdots, \bar{x}_N) \in T^N, \quad \widetilde{x} = (\widetilde{x}_1, \widetilde{x}_2, \cdots, \widetilde{x}_N) \in \widetilde{E}.$$

Since the matrix $A(q)$ defined by (1.3) is positive definite and periodic in q, there exists a constant $\lambda > 0$ depending only on masses and lengths such that

$$\lambda = \inf\left\{ \frac{A(q)p\cdot p}{|p|^2} \mid q \in T^N, p \in \mathbb{R}^N \setminus \{0\} \right\}. \tag{2.3}$$

Let

$$\gamma_1 = 2\pi^2 \left(\sum_{i=1}^{N} \alpha_i \ell_i^2 v_i^2 + \sum_{i<j, v_i = v_j} 2\alpha_j \ell_i \ell_j v_i v_j \right). \tag{2.4}$$

We have

Proposition 2.1.

$$\mathscr{L}_1(x) \geqslant \frac{\lambda}{2}||\dot{x}||^2 + T^{-1}2\pi^2|v|^2\lambda, \tag{2.5}$$

$$\mathscr{L}_1(\bar{x}) \leqslant T^{-1}\gamma_1,\ \mathscr{L}_2(\bar{x}) = T\sum_{i\in I_0} \beta_i \cos\bar{x}_i,\ \mathscr{L}_3(\bar{x}) = \int_0^T f(t)\cdot\frac{2\pi v t}{T}\mathrm{d}t = f_v, \tag{2.6}$$

$$|\mathscr{L}_2(x) - \mathscr{L}_2(\bar{x})| \leqslant \frac{T^{\frac{3}{2}}}{2\pi}\sum_{i=1}^{N} \beta_i ||\dot{x}_i||, \tag{2.7}$$

$$|\mathscr{L}_3(x) - \mathscr{L}_3(\bar{x})| \leqslant \frac{T}{2\pi}\sum_{i=1}^{N} ||f||\cdot||\dot{x}_i|| \leqslant \frac{T^{\frac{3}{2}}}{2\pi}\sum_{i=1}^{N} M_0||\dot{x}_i||. \tag{2.8}$$

Proof. For (2.5), by (2.2) and (2.3), we have

$$\mathscr{L}_1(x) = \frac{1}{2}\langle A(q)\dot{q}, \dot{q}\rangle_{L^2} \geqslant \frac{\lambda}{2}\left|\left|\dot{x} + \frac{2\pi v}{T}\right|\right|^2 = \frac{\lambda}{2}||\dot{x}||^2 + \frac{2\pi^2|v|^2\lambda}{T}.$$

For (2.6), we claim that

$$\int_0^T \cos\left(\bar{x}_i + 2\pi v_i \frac{t}{T}\right)\mathrm{d}t = \begin{cases} 0, & v_i \neq 0, \\ T\cos\bar{x}_i, & v_i = 0, \end{cases} \quad 1 \leqslant i \leqslant N, \tag{2.9}$$

$$\left| \int_0^T \cos\left(x_i + 2\pi v_i \frac{t}{T}\right)\mathrm{d}t - \int_0^T \cos\left(\bar{x}_i + 2\pi v_i \frac{t}{T}\right)\mathrm{d}t \right| \leqslant \frac{T\sqrt{T}}{2\pi}||\dot{x}_i||. \tag{2.10}$$

In fact

$$\begin{aligned}
&\left|\int_0^T \cos\left(x_i+2\pi v_i\frac{t}{T}\right)\mathrm{d}t-\int_0^T \cos\left(\bar{x}_i+2\pi v_i\frac{t}{T}\right)\mathrm{d}t\right|\\
&=\left|\int_0^T \left(\cos\left(x_i+2\pi v_i\frac{t}{T}\right)-\cos\left(\bar{x}_i+2\pi v_i\frac{t}{T}\right)\right)\mathrm{d}t\right|\\
&=\left|-2\int_0^T \sin\left(\frac{\widetilde{x}_i}{2}+\bar{x}_i+2\pi v_i\frac{t}{T}\right)\sin\left(\frac{\widetilde{x}_i}{2}\right)\mathrm{d}t\right|\\
&\leqslant 2\left(\int_0^T \sin^2\left(\frac{\widetilde{x}_i}{2}+\bar{x}_i+2\pi v_i\frac{t}{T}\right)\mathrm{d}t\right)^{\frac{1}{2}}\left(\int_0^T \sin^2\frac{\widetilde{x}_i}{2}\mathrm{d}t\right)^{\frac{1}{2}}\\
&\leqslant 2\sqrt{T}\left(\int_0^T \sin^2\frac{\widetilde{x}_i}{2}\mathrm{d}t\right)^{\frac{1}{2}}\leqslant 2\sqrt{T}\left(\int_0^T \frac{\widetilde{x}_i^2}{4}\mathrm{d}t\right)^{\frac{1}{2}}\\
&=\sqrt{T}\|\widetilde{x}_i\|\leqslant\frac{T\sqrt{T}}{2\pi}\|\dot{x}_i\|.
\end{aligned}$$

The last inequality follows from the Wirtinger's inequality.

By (2.2), (1.3), (1.7) and (2.9), we have

$$\begin{aligned}
\mathscr{L}_1(\bar{x})&=\sum_{i=1}^N\alpha_i\ell_i^2\cdot 2\pi^2v_i^2\frac{1}{T}+\sum_{1\leqslant i<j\leqslant N}\alpha_j\ell_i\ell_j\cdot\frac{4\pi^2}{T^2}v_iv_j\int_0^T\cos\left(\bar{x}_i-\bar{x}_j+2\pi(v_i-v_j)\frac{t}{T}\right)\mathrm{d}t\\
&=\frac{2\pi^2}{T}\sum_{i=1}^N\alpha_i\ell_i^2v_i^2+\sum_{i<j,v_i=v_j}\alpha_j\ell_i\ell_j\cdot\frac{4\pi^2}{T^2}v_iv_j\int_0^T\cos(\bar{x}_i-\bar{x}_j)\mathrm{d}t\\
&\leqslant\frac{2\pi^2}{T}\sum_{i=1}^N\alpha_i\ell_i^2v_i^2+\frac{4\pi^2}{T}\sum_{i<j,v_i=v_j}\alpha_j\ell_i\ell_jv_iv_j=T^{-1}\gamma_1.
\end{aligned}$$

By (2.2), (1.7), (1.4) and (1.12), we have

$$\begin{aligned}
\mathscr{L}_2(\bar{x})&=\int_0^T V\left(\bar{x}+2\pi v\frac{t}{T}\right)\mathrm{d}t\\
&=\int_0^T\sum_{i=1}^N\beta_i\cos\left(\bar{x}_i+2\pi v_i\frac{t}{T}\right)\mathrm{d}t=T\sum_{i\in I_0}\beta_i\cos\bar{x}_i,\\
\mathscr{L}_3(\bar{x})&=\int_0^T f(t)\cdot\left(\bar{x}+2\pi v\frac{t}{T}\right)\mathrm{d}t=\int_0^T f(t)\cdot\frac{2\pi vt}{T}\mathrm{d}t.
\end{aligned}$$

The inequality (2.7) follows from (2.2), (1.4) and (2.10).

For (2.8), by (2.2), (1.7) and (1.12), we have

$$\begin{aligned}
|\mathscr{L}_3(x)-\mathscr{L}_3(\bar{x})|&=\left|\int_0^T f\cdot q\,\mathrm{d}t-\int_0^T f2\pi v\frac{t}{T}\mathrm{d}t\right|=\left|\int_0^T f\cdot(\bar{x}+\widetilde{x})\mathrm{d}t\right|\\
&=\left|\int_0^T f\cdot\widetilde{x}\mathrm{d}t\right|=\left|\int_0^T\sum_{i=1}^N f_i\cdot\widetilde{x}_i\mathrm{d}t\right|\leqslant\sum_{i=1}^N\int_0^T|f_i|\cdot|\widetilde{x}_i|\mathrm{d}t\\
&\leqslant\sum_{i=1}^N\|f_i\|\cdot\|\widetilde{x}_i\|\leqslant\sum_{i=1}^N\|f_i\|\cdot\frac{T}{2\pi}\|\dot{x}_i\|\leqslant\frac{T\|f\|}{2\pi}\|\dot{x}\|.
\end{aligned}$$

The third equality follows from (2.6), and the last inequality follows from the Wirtinger's inequality. ∎

By the above Proposition, we have

$$\left|\mathscr{L}_2(x)+\mathscr{L}_3(x)-\left(\mathscr{L}_2(\bar{x})+\mathscr{L}_3(\bar{x})\right)\right| \leqslant \frac{T^{\frac{3}{2}}}{2\pi}\sum_{i=1}^{N}(\beta_i+M_0)\|\dot{x}_i\|. \tag{2.11}$$

Proposition 2.2. *For the functional* $\mathscr{L}$ *defined by* (1.8)*, we have*
(i) $\mathscr{L}$ *is bounded from below.*
(ii) $\mathscr{L}$ *satisfies the (PS) condition.*

Proof. For (i), by , we have

$$\begin{aligned}\mathscr{L}(x) &\geqslant \frac{\lambda}{2}\|\dot{x}\|^2+T^{-1}2\pi^2|v|^2\lambda-T\sum_{i=1}^{N}\beta_i-\frac{T^{\frac{3}{2}}\|f\|}{2\pi}\|\dot{x}\|+f_v\\ &=\frac{\lambda}{2}\sum_{i=1}^{N}\left(\|\dot{x}\|-\frac{T^{\frac{3}{2}}\|f\|}{2\pi\lambda}\right)^2+a_0,\ \forall x\in\mathscr{E},\end{aligned} \tag{2.12}$$

where

$$a_0=T^{-1}2\pi^2|v|^2\lambda+f_v-T\sum_{i=1}^{N}\beta_i-\frac{T^3\|f\|^2}{8\pi^2\lambda}. \tag{2.13}$$

For (ii), let $\{x_m\}$ be a sequence in $\mathscr{E}$ such that

$$\mathscr{L}(x_m)\leqslant C \text{ and } \mathscr{L}'(x_m)\to 0 \text{ as } m\to\infty. \tag{2.14}$$

Then (2.12) yields that

$$\|\dot{x}_m\|\leqslant\|x_m\|_E\leqslant C', \tag{2.15}$$

i.e., $\{x_m\}$ is bounded in $\mathscr{E}$ since T^N is compact. Without loss of generality, assume that $x_m\rightharpoonup x_0\in\mathscr{E}$ and $x_m\to x_0\in C(S^1,\mathbb{R}^N)$ as $m\to\infty$, i.e.,

$$\lim_{m\to\infty}\|x_m-x_0\|_{C^0}=0. \tag{2.16}$$

Note that for arbitrary $x,y\in\mathscr{E}$, by using the substitution (1.7) for simplicity, we have

$$\mathscr{L}'(x)y=\mathscr{L}_1'(x)y+\langle V'(q)+f,y\rangle, \tag{2.17}$$

where

$$\begin{aligned}\mathscr{L}_1'(x)y&=\langle A(q)\dot{q},\dot{y}\rangle+\left\langle\left(A'(q)y\right)\dot{q},\dot{q}\right\rangle,\\ \left\langle\left(A'(q)y\right)\dot{q},\dot{q}\right\rangle&=\int_0^T\sum\alpha_j\ell_i\ell_j\left(-\sin(q_i-q_j)(y_i-y_j)\right)\dot{q}_i\dot{q}_j\mathrm{d}t.\end{aligned}$$

Then

$$\begin{aligned}\mathscr{L}_1'(x+y)y-\mathscr{L}_1'(x)y&=\langle A(q+y)(\dot{q}+\dot{y}),\dot{y}\rangle+\left\langle(A'(q+y)y)(\dot{q}+\dot{y}),\dot{q}+\dot{y}\right\rangle\\ &\quad-\langle A(q)\dot{q},\dot{y}\rangle-\left\langle\left(A'(q)y\right)\dot{q},\dot{q}\right\rangle\\ &=\langle A(q+y)\dot{y},\dot{y}\rangle+\left\langle\left(A(q+y)-A(q)\right)\dot{q},\dot{y}\right\rangle\\ &\quad+\left\langle\left(A'(q+y)y-A'(q)y\right)\dot{q},\dot{q}\right\rangle\\ &\quad+2\left\langle\left(A'(q+y)y\right)\dot{q},\dot{y}\right\rangle+\left\langle\left(A'(q+y)y\right)\dot{y},\dot{y}\right\rangle\\ &\geqslant\lambda\|\dot{y}\|^2+o\left(\|y\|\right).\end{aligned}$$

We have

$$\mathscr{L}_1'(x_m)(x_m - x_0) - \mathscr{L}_1'(x_0)(x_m - x_0) \geqslant \lambda ||\dot{x}_m - \dot{x}_0||^2 + o(||x_m - x_0||). \tag{2.18}$$

By (2.17) and (2.18), we have

$$\begin{aligned}
&\mathscr{L}'(x_m)(x_m - x_0) - \mathscr{L}'(x_0)(x_m - x_0)\\
&= \mathscr{L}_1'(x_m)(x_m - x_0) + \langle V'(q_m) + f, x_m - x_0\rangle\\
&\ - \mathscr{L}_1'(x_0)(x_m - x_0) - \langle V'(q_0) + f, x_m - x_0\rangle\\
&\geqslant \lambda ||\dot{x}_m - \dot{x}_0||^2 + o(||x_m - x_0||).
\end{aligned}$$

Then (2.14) and (2.16) yield $||\dot{x}_m - \dot{x}_0|| \to 0$ as $m \to \infty$. Hence $x_m \to x_0$ in $\mathscr{E}$. ∎

Proof of Theorem 1. For (i), by Proposition 2.2 and Ljusternik-Schnirelman theory, we have

$$^{\#}\mathscr{P}(v,T) \geqslant \mathrm{cat}(T^N) = N + 1.$$

If all solutions are nondegenerate, by Morse inequalities, we have

$$^{\#}\mathscr{P}(v,T) \geqslant \text{ the sum of Betti numbers of } T^N = 2^N.$$

For (ii), by Proposition 2.2 and S^1-equivariant Ljusternik-Schnirelman theory, we have

$$^{\#}\mathscr{P}(v,T) \geqslant \mathrm{cat}(T^N/S^1) = \mathrm{cat}(T^{N-1}) = N.$$

If all solutions are nondegenerate, by Morse inequalities, we have

$$^{\#}\mathscr{P}(v,T) \geqslant \text{ the sum of Betti numbers of } T^{N-1} = 2^{N-1}. \blacksquare$$

3 An abstract critical point result

In [4], K. C. Chang, Y. Long and E. Zehnder prove an abstract critical point result (Theorem 3.1) and apply it to oscillatory solutions of (1.5). In this section, we establish corresponding result and apply it to rotational solutions. The difference is that N_0 defined by (1.13) satisfies

$$N_0 \begin{cases} = N, & \text{oscillatory case, i.e.,} v = 0,\\ \leqslant N - 1, & \text{rotational case, i.e.,} v \neq 0.\end{cases}$$

The main idea is to find an m-dimensonal subtorus $T^m \subset T^N$ with an $(m-1)$-dimensonal subtorus $T^{m-1} \subset T^m$ such that the inclusion

$$(T^m, T^{m-1}) \hookrightarrow (I^b, I^a)$$

induces an injective map in homology and a surjective map in cohomology. By Ljusternik-Schirelman theory, there exist at least m critical points between the two levels a and b.



Let

$$n = 2^{N_0}. \tag{3.1}$$

For each $1 \leqslant k \leqslant n$, represent $k-1$ by binary numbers as follows

$$k-1 = \sum_{i=1}^{N_0} \tau_i(k) 2^{N_0-i},\ \tau_i(k) \in \{0,1\}, \tag{3.2}$$

and set

$$z(k) = \big(z_1(\tau_1(k)), z_2(\tau_2(k)), \cdots, z_{N_0}(\tau_{N_0}(k))\big) \in T^{N_0}, \tag{3.3}$$

where $z_i : \{0,1\} \to S^1,\ 1 \leqslant i \leqslant N_0$, are injective maps.

Define an $(N-N_0)$-dimensional subtorus $S_k \subset T^N$ by

$$S_k = \{z(k)\} \times T^{N-N_0}. \tag{3.4}$$

For k, j satisfying $1 \leqslant j \leqslant k$ and

$$k - j = 2^{N_0-m},\ \text{for some } 1 \leqslant m \leqslant N_0,$$

define an $(N-N_0+1)$-dimensional subtorus $T_{k,j} \subset T^N$ containing subtorus S_k and S_j as follows

$$T_{k,j} = \big\{(x_1, \cdots, x_{N_0}) \in T^{N_0} \ \big|\ x_i = z_i(\tau_i(k)),\ i \neq m\big\} \times T^{N-N_0} \cong T^{N-N_0+1}. \tag{3.5}$$

For $1 \leqslant k \leqslant n,\ 1 \leqslant i \leqslant N_0$, let

$$T_{k,i}^{N-1} = \big\{(y_1, \cdots, y_N) \ \big|\ y_i = z_i(1-\tau_i(k)) \text{ and } y_j \in S^1 \text{ for } j \neq i\big\} \cong T^{N-1},$$
$$Q_k = \bigcup_{i=1}^{N_0} T_{k,i}^{N-1},\ M_k = \bigcap_{j=k}^{n} Q_j. \tag{3.6}$$

By the proof of Lemma 3.1 of [4], there is a strong deformation retraction

$$r_k : T^N \setminus S_k \to Q_k. \tag{3.7}$$

Let

$$W_k = T^N \setminus \left[S_n \cup \left(\bigcup_{i=k}^{n-1} r_n^{-1} \cdots r_{i+1}^{-1}(S_i) \right) \right], \tag{3.8}$$

$$R_k = r_k \circ r_{k+1} \circ \cdots \circ r_n : \mathrm{dom}(r_k \circ r_{k+1} \circ \cdots \circ r_n) = W_k \to T^N. \tag{3.9}$$

Proposition 3.1. *Assume $I \in C^1(\mathscr{E}, \mathbb{R})$ satisfies (PS) condition and $a_0 = \inf\limits_{x \in \mathscr{E}} I(x) > -\infty$. If there exist real numbers $a_0 < a_1 < \cdots < a_n$, where $n = 2^{N_0}$ such that*

$$I_{T^N \times \{0\}} < a_n,\ I|_{M_{k+1} \times \{0\}} < a_k < I|_{p^{-1} R_{k+2}^{-1}(S_{k+1})},\ 1 \leqslant k \leqslant n-1, \tag{3.10}$$

(Here we set $R_{n+1}^{-1}(S_n) = S_n$). Denote by Crit(I) the set of critical points of I, we have

$$^{\#}Crit(I) \geqslant (N - N_0 + 1)2^{N_0}.$$

If I is S^1-equivariant, then

$$^{\#}Crit(I) \geqslant (N - N_0)2^{N_0}.$$

Proof. For brevity, let

$$m = N - N_0 + 1.$$

We claim that

$$^{\#}\Big(\mathrm{Crit}(I) \cap \mathring{I}^{a_1}\Big) \geqslant m, \quad k = 1, \tag{3.11}$$

$$^{\#}\Big(\mathrm{Crit}(I) \cap \big(\mathring{I}^{a_k} \setminus \mathring{I}^{a_{k-1}}\big)\Big) \geqslant m, \quad 2 \leqslant k \leqslant n-1, \tag{3.12}$$

$$^{\#}\Big(\mathrm{Crit}(I) \cap \big(\mathscr{E} \setminus I^{a_{n-1}}\big)\Big) \geqslant m, \quad k = n, \tag{3.13}$$

where

$$I^a = \{x \in \mathscr{E} \mid I(x) \leqslant a\} \quad \text{and} \quad \mathring{I}^a = \{x \in \mathscr{E} \mid I(x) < a\}.$$

By (3.11)-(3.13), we have

$$^{\#}\mathrm{Crit}(I) \geqslant m + (n-2)m + m = mn = (N - N_0 + 1)2^{N_0}.$$

Case 1. $k = 1$. By (3.15), there exists $\delta_1 \in (0, a_1 - a_0)$ such that

$$S_1 \times \{0\} \subset I^{a_1 - \delta_1}.$$

The Ljusternik-Schnirelman category of $I^{a_1 - \delta_1}$ is

$$\mathrm{cat}(I^{a_1 - \delta_1}) \geqslant \mathrm{cat}(S_1) = \mathrm{cat}(T^{m-1}) = m.$$

As $a_0 = \inf I > -\infty$ and I satisfies (PS), we conclude (3.11).
Case 2. $2 \leqslant k \leqslant n-1$. We proceed in several steps.

Step 1. Choose $\delta \in (0, a_k - a_{k-1})$ sufficiently small, abbreviate

$$b = a_k - \delta,\ a = a_{k-1}$$

and set

$$\begin{aligned} X &= \mathscr{E} \setminus \left[p^{-1}(S_n) \cup p^{-1}R_n^{-1}(S_{n-1}) \cup \cdots \cup p^{-1}R_{k+2}^{-1}(S_{k+1})\right], \\ X' &= \mathscr{E} \setminus \left[p^{-1}(S_n) \cup p^{-1}R_n^{-1}(S_{n-1}) \cup \cdots \cup p^{-1}R_{k+1}^{-1}(S_k)\right], \\ Y &= I^b,\ Y' = I^a,\ Z = M_{k+1} \times \{0\},\ Z' = M_k \times \{0\}. \end{aligned}$$

By (3.15), we have

$$\begin{array}{ccccc} X & \supset & Y & \supset & Z \\ \cup & & \cup & & \cup \\ X' & \supset & Y' & \supset & Z'. \end{array}$$

Step 2. Consider the following inclusion maps

$$(T_{k,k'}, S_{k'}) \xrightarrow{\rho} (M_{k+1}, M_k) \xrightarrow[x \mapsto (x,0)]{\beta} (Z, Z') \xrightarrow{j} (Y, Y').$$

By the proof on page 195 of [4], the continuous map

$$\psi = j \circ \beta \circ \rho : (T_{k,k'}, S_{k'}) \to (I^b, I^a)$$

induces a monomorphism

$$\psi_* : H_*(T_{k,k'}, S_{k'}) \to H_*(I^b, I^a),$$

and an epimorphism

$$\psi^* : H^*(I^b) \to H^*(T_{k,k'}).$$

Step 3. We claim that

$$H_q(T_{k,k'}, S_{k'}) \cong H_q\left(T^m, T^{m-1}\right) \neq 0,\ 1 \leqslant q \leqslant m.$$

In fact, the first isomorphism follows from (3.4) and (3.5). Consider the following long exact sequence

$$\cdots \to H_q(T^{m-1}) \xrightarrow{i_q} H_q(T^m) \xrightarrow{j_q} H_q(T^m, T^{m-1}) \xrightarrow{\partial_q} H_{q-1}(T^{m-1}) \xrightarrow{i_{q-1}} \cdots$$

We have

$$\begin{aligned} H_q(T^{m-1}) &\cong \mathrm{Ker} i_q \oplus \mathrm{Im} i_q \cong \mathbb{Z}^{C^q_{m-1}}, \\ H_q(T^m) &\cong \mathrm{Ker} j_q \oplus \mathrm{Im} j_q \cong \mathbb{Z}^{C^q_m}, \\ H_q(T^m, T^{m-1}) &\cong \mathrm{Ker}\partial_q \oplus \mathrm{Im}\partial_q, \\ \mathrm{Im} i_q = \mathrm{Ker} j_q,\ &\mathrm{Im} j_q = \mathrm{Ker}\partial_q,\ \mathrm{Im}\partial_q = \mathrm{Ker} i_{q-1}. \end{aligned}$$

Hence, for $1 \leqslant q \leqslant m$, the q-th Betti-number of the pair (T^m, T^{m-1}) satisfies

$$\begin{aligned} b_q(T^m, T^{m-1}) &= \dim H_q(T^m, T^{m-1}) \\ &= C^q_m - C^q_{m-1} + \dim \mathrm{Ker} i_q + \dim \mathrm{Ker} i_{q-1} \\ &\geqslant C^q_m - C^q_{m-1} \geqslant 1. \end{aligned}$$

Step 4. Choose homology classes

$$0 \neq [u_q] \in H_q(T_{k,k'}, S_{k'}),\ 1 \leqslant q \leqslant m,$$

and a cohomology class $0 \neq \theta \in H^1(T_{k,k'}) \cong H^1(T^m)$ such that

$$[u_{q+1}] \cap \theta = [u_q],\ 1 \leqslant q \leqslant m-1.$$

Let

$$[\sigma_q] = \psi_*[u_q],\ 1 \leqslant q \leqslant m,$$

and choose

$$\omega \in (\psi^*)^{-1}(\theta),$$

then

$$0 \neq [\sigma_q] \in H_q(I^b, I^a) \text{ and } 0 \neq \omega \in H^1(I^b).$$

By Proposition 5.6.16 (on page 254) of [16], we have

$$\begin{aligned}[\sigma_{q+1}] \cap \omega &= \left(\psi_*([u_{q+1}])\right) \cap \omega = \psi_*([u_{q+1} \cap \psi^*\omega]) \\ &= \psi_*([u_{q+1} \cap \theta]) = \psi_*([u_q]) = [\sigma_q],\ 1 \leqslant q \leqslant m-1.\end{aligned}$$

By definition 1.1 (on page 10) of [5], we have

$$[\sigma_1] < [\sigma_2] < \cdots < [\sigma_m].$$

Apply the Ljusternik-Schnirelman theory (see Corollary 3.3 (on page 106) of [5]), we conclude (3.12).

Case 3. $k = n$. The proof is similar to that of Case 1. In Step 1, we set

$$X = Y = \mathscr{E},\ Z = T^N \times \{0\},\ Z' = M_n \times \{0\} = Q_n \times \{0\},$$

and conclude $N - N_0 + 1$ critical points above the level a_{n-1}.

If I is S^1-equivariant, then replace X, Y, Z, X', Y', Z' by their quotient spaces under the S^1-action. We have

$$\begin{aligned}\operatorname{cat}(I^{a_1-\delta_1}/S^1) &\geqslant \operatorname{cat}(S_1/S^1) = \operatorname{cat}(T^{m-2}) = m-1, \\ H_q(T_{k,k'}/S^1, S_{k'}/S^1) &\cong H_q\left(T^m/S^1, T^{m-1}/S^1\right) \\ &\cong H_q\left(T^{m-1}, T^{m-2}\right) \neq 0,\ 1 \leqslant q \leqslant m-1.\end{aligned}$$

Then repeat Step 4 in Case 2 by replacing m by $m-1$. ∎

For each homeomorphism

$$\phi : T^N \to T^N, \tag{3.14}$$

set

$$\begin{gathered}\widetilde{S}_k = \phi^{-1}(S_k) \cong T^{N-N_0},\ \widetilde{T}_{k,j} = \phi^{-1}\left(T_{k,j}\right) \cong T^{N-N_0+1}, \\ \widetilde{T}^{N-1}_{k,i} = \phi^{-1}\left(T^{N-1}_{k,i}\right) \cong T^{N-1},\ \widetilde{Q}_k = \phi^{-1}(Q_k),\ \widetilde{M}_k = \phi^{-1}(M_k), \\ \widetilde{r}_k = \phi^{-1} \circ r_k,\ \widetilde{W}_k = \phi^{-1}(W_k), \widetilde{R}_k = \phi^{-1} \circ (R_k) = \widetilde{r}_k \circ \widetilde{r}_{k+1} \circ \cdots \circ \widetilde{r}_n : \widetilde{W}_k \to T^N.\end{gathered}$$

By (3.7), the map

$$\widetilde{r}_k : T^N \setminus \widetilde{S}_k \to \widetilde{Q}_k,$$

is a strong deformation retraction. By (3.6), we have

$$\widetilde{Q}_k = \bigcup_{i=1}^{N_0} T^{N-1}_{k,i},\ \widetilde{M}_k = \bigcap_{j=k}^{n} \widetilde{Q}_j.$$

Since the above homeomorphism ϕ induces isomorphisms in homology and cohomology, by Proposition 3.1, we have

Corollary 1. *Assume $I \in C^1(\mathscr{E},\mathbb{R})$ satisfies (PS) condition and $a_0 = \inf_{x\in\mathscr{E}} I(x) > -\infty$. If there exist real numbers $a_0 < a_1 < \cdots < a_n$, where $n = 2^{N_0}$ such that*

$$I_{T^N\times\{0\}} < a_n,\ I|_{\widetilde{M}_{k+1}\times\{0\}} < a_k < I|_{p^{-1}\widetilde{R}^{-1}_{k+2}(\widetilde{S}_{k+1})},\ 1 \leqslant k \leqslant n-1, \tag{3.15}$$

(Here we set $R^{-1}_{n+1}(S_n) = S_n$). Denote by Crit(I) the set of critical points of I, we have

$$^{\#}Crit(I) \geqslant (N - N_0 + 1)2^{N_0}.$$

If ϕ and I are S^1-equivariant, then

$$^{\#}Crit(I) \geqslant (N - N_0)2^{N_0}.$$

In (3.3), we set

$$z(k) = \big(\pi(1-\tau_1(k)), \pi(1-\tau_2(k)), \cdots, \pi(1-\tau_{N_0}(k))\big). \tag{3.16}$$

The following result will be utilized in the proof of Theorem 2.

Proposition 3.2. *For $2 \leqslant k \leqslant 2^{N_0} - 1$, let*

$$O_k = \left\{ (y_1, y_2, \cdots, y_N) \in T^N \;\middle|\; y_i = 0,\ \text{if } \tau_i(k) = 1 \right\}. \tag{3.17}$$

Then

$$S_k \subset R^{-1}_{k+1}(S_k) \subset O_k. \tag{3.18}$$

Proof. For induction on N_0 and brevity, let $m = N_0$,

$$\begin{aligned} &z^{(m)}(k) = z(k), \quad S^{(m)}_k = S_k, \quad O^{(m)}_k = O_k, \\ &Q^{(m)}_k = Q_k, \quad M^{(m)}_k = M_k, \quad r_{m,k} = r_k, \quad R_{m,k+1} = R_{k+1}. \end{aligned}$$

The map $r_{m,k} = r_k : T^N \setminus S^{(m)}_k \to Q^{(m)}_k$ is defined by

$$\left(y^{(m)}, y'\right) \mapsto \left(\sigma y^{(m)} + (1-\sigma) z^{(m)}(k), y'\right),$$

where $\sigma \geqslant 1$ is uniquely determined by k and $y^{(m)}$ (c.f. Lemma 3.1 of [4]). We have

$$r^{-1}_{m,k}\left(w^{(m)}, \omega_{m+1}\right) = \left\{ \lambda w^{(m)} + (1-\lambda) z^{(m)}(k) \;\middle|\; 0 < \lambda \leqslant 1 \right\} \times \{\omega_{m+1}\}, \tag{3.19}$$

$$r_{m,s+1}^{-1} r_{m,s}^{-1}\left(O_k^{(m)}\right) = r_{m,s+1}^{-1}\big(r_{m,s}^{-1}\left(O_k^{(m)}\right) \cap Q_{s+1}^{(m)}\big),\ k+1 \leqslant s \leqslant 2^m, \tag{3.20}$$

$$\begin{aligned} r_{m,q}^{-1} \cdots r_{m,p}^{-1}(\omega) \subset \Bigg\{ & \lambda_q \cdots \lambda_p \omega + \sum_{i=p}^{q-1} \lambda_q \cdots \lambda_{i+1}(1-\lambda_i) z^{(m)}(i) \\ & +(1-\lambda_q) z^{(m)}(q) \,\Bigg|\, 0 < \lambda_i \leqslant 1 \Bigg\} \times T^{N-m}. \end{aligned} \tag{3.21}$$

Note that $R_{m,k+1}\left(S_k^{(m)}\right) = S_k^{(m)}$. We claim that

$$R_{m,s}^{-1}\left(O_k^{(m)}\right) \subset O_k^{(m)},\ k \leqslant 2^m - 1,\ s = 1,\ k+1. \tag{3.22}$$

	$1 \leqslant k \leqslant 2^{m-1}$	$2^{m-1}+1 \leqslant k \leqslant 2^m$
$z^{(m)}(k)$	$\left(\pi, z^{(m-1)}(k)\right)$	$\left(0, z^{(m-1)}\left(k-2^{m-1}\right)\right)$
$S_k^{(m)}$	$\{\pi\} \times S_k^{(m-1)}$	$\{0\} \times S_{k-2^{m-1}}^{(m-1)}$
$O_k^{(m)}$	$S^1 \times O_k^{(m-1)}$	$\{0\} \times O_{k-2^{m-1}}^{(m-1)}$
$Q_k^{(m)}$	$\left(\{0\} \times T^{N-1}\right) \bigcup \left(S^1 \times Q_k^{(m-1)}\right)$	$\left(\{\pi\} \times T^{N-1}\right) \bigcup \left(S^1 \times Q_{k-2^{m-1}}^{(m-1)}\right)$

If $m = 2$, we have $2 \leqslant k \leqslant 2^2 - 1 = 3$,

$$\begin{aligned} O_2^{(2)} &= S^1 \times \{0\} \times T^{N-2}, \\ O_3^{(2)} &= \{0\} \times S^1 \times T^{N-2}. \end{aligned}$$

Then

$$\begin{aligned} R_{2,3}^{-1}\left(O_2^{(2)}\right) &= r_{2,4}^{-1} r_{2,3}^{-1}\left(O_2^{(2)}\right) = r_{2,4}^{-1}\left(r_{2,3}^{-1}\left(O_2^{(2)}\right) \cap Q_4^{(2)}\right) \subset r_{2,4}^{-1}\left(O_2^{(2)}\right) \subset O_2^{(2)}, \\ R_{2,1}^{-1}\left(O_2^{(2)}\right) &= r_{2,4}^{-1} r_{2,3}^{-1} r_{2,2}^{-1} r_{2,1}^{-1}\left(O_2^{(2)}\right) = r_{2,4}^{-1} r_{2,3}^{-1} r_{2,2}^{-1}\left(r_{2,1}^{-1}\left(O_2^{(2)}\right) \cap Q_2^{(2)}\right) \\ &\subset r_{2,4}^{-1} r_{2,3}^{-1} r_{2,2}^{-1}\left(O_2^{(2)}\right) = r_{2,4}^{-1} r_{2,3}^{-1}\left(r_{2,2}^{-1}\left(O_2^{(2)}\right) \cap Q_3^{(2)}\right) \\ &\subset r_{2,4}^{-1} r_{2,3}^{-1}\left(O_2^{(2)}\right) \subset O_2^{(2)}, \\ R_{2,4}^{-1}\left(O_3^{(2)}\right) &= r_{2,4}^{-1}\left(O_3^{(2)}\right) \subset O_3^{(2)}, \quad R_{2,1}^{-1}\left(O_3^{(2)}\right) = r_{2,4}^{-1} r_{2,3}^{-1} r_{2,2}^{-1} r_{2,1}^{-1}\left(O_3^{(2)}\right) = \emptyset. \end{aligned}$$

Assume (3.22) holds for $m = p$. We carry out the proof for $m = p+1$ as follows.
Case 1. $2 \leqslant k \leqslant 2^{m-1} - 1$. We have $O_k^{(m)} = S^1 \times O_k^{(m-1)}$. Then

$$\begin{aligned} r_{m,2^{m-1}}^{-1} \cdots r_{m,s}^{-1}\left(O_k^{(m)}\right) &\subset S^1 \times R_{m-1,s}^{-1}\left(O_k^{(m-1)}\right) \subset S^1 \times O_k^{(m-1)},\ s = 1,\ k+1, \\ R_{m,s}^{-1}\left(O_k^{(m)}\right) &= R_{m,2^{m-1}+1}^{-1}\left(r_{m,2^{m-1}}^{-1} \cdots r_{m,s}^{-1}\left(O_k^{(m)}\right)\right) \\ &\subset R_{m,2^{m-1}+1}^{-1}\left(S^1 \times O_k^{(m-1)}\right) \\ &\subset S^1 \times \left(O_k^{(m-1)} \cup R_{m-1,1}^{-1}\left(O_k^{(m-1)}\right)\right) \subset O_k^{(m)}. \end{aligned}$$

Case 2. $k = 2^{m-1}$. We have $O^{(m)}_{2^{m-1}} = S^1 \times O^{(m-1)}_{2^{m-1}}$. Then

$$\begin{aligned}
R^{-1}_{m,2^{m-1}+1}\left(O^{(m)}_{2^{m-1}}\right) &\subset S^1 \times \left(O^{(m-1)}_{2^{m-1}} \cup R^{-1}_{m-1,1}\left(O^{(m-1)}_{2^{m-1}}\right)\right) \subset O^{(m)}_{2^{m-1}}, \\
R^{-1}_{m,1}\left(O^{(m)}_{2^{m-1}}\right) &= R^{-1}_{m,2^{m-1}+1}\left(r^{-1}_{m,2^{m-1}} \cdots r^{-1}_{m,1}\left(O^{(m)}_{2^{m-1}}\right)\right) \\
&\subset R^{-1}_{m,2^{m-1}+1}\left(S^1 \times R^{-1}_{m-1,1}\left(O^{(m-1)}_{2^{m-1}}\right)\right) \\
&\subset R^{-1}_{m,2^{m-1}+1}\left(S^1 \times O^{(m-1)}_{2^{m-1}}\right) \\
&\subset S^1 \times \left(O^{(m-1)}_{2^{m-1}} \cup R^{-1}_{m-1,1}\left(O^{(m-1)}_{2^{m-1}}\right)\right) \subset O^{(m)}_{2^{m-1}}.
\end{aligned}$$

Case 3. $2^{m-1} + 1 \leqslant k \leqslant 2^m - 1$. We have $1 \leqslant k - 2^{m-1} \leqslant 2^{m-1} - 1$, $O^{(m)}_k = \{0\} \times O^{(m-1)}_{k-2^{m-1}}$ and

$$r^{-1}_{m,s}\left(O^{(m)}_k\right) = \{0\} \times r^{-1}_{m-1,s-2^{m-1}}\left(O^{(m-1)}_{k-2^{m-1}}\right),\ k+1 \leqslant s \leqslant 2^m.$$

Then

$$\begin{aligned}
R^{-1}_{m,k+1}\left(O^{(m)}_k\right) &= r^{-1}_{m,2^m} \cdots r^{-1}_{m,k+1}\left(O^{(m)}_k\right) \\
&= \{0\} \times R^{-1}_{m-1,k-2^{m-1}+1}\left(O^{(m-1)}_{k-2^{m-1}}\right) \\
&\subset \{0\} \times O^{(m-1)}_{k-2^{m-1}} = O^{(m)}_k, \\
R^{-1}_{m,1}\left(O^{(m)}_k\right) &= R^{-1}_{m,2^{m-1}+1}\left(r^{-1}_{m,2^{m-1}} \cdots r^{-1}_{m,1}\left(O^{(m)}_k\right)\right) \\
&= R^{-1}_{m,2^{m-1}+1}\left(\{0\} \times R^{-1}_{m-1,1}\left(O^{(m-1)}_{k-2^{m-1}}\right)\right) \\
&\subset R^{-1}_{m,2^{m-1}+1}\left(\{0\} \times O^{(m-1)}_{k-2^{m-1}}\right) \\
&= \{0\} \times R^{-1}_{m-1,1}\left(O^{(m-1)}_{k-2^{m-1}}\right) \subset \{0\} \times O^{(m-1)}_{k-2^{m-1}} \subset O^{(m)}_k. \blacksquare
\end{aligned}$$

4 Proof of Theorem 2

4.1 Restrictions of masses, lengths and periods

Choose a permutation σ of $\{1, 2, \cdots, N\}$ such that

$$\begin{gathered}
I_0 = \{\sigma(1),\ \sigma(2),\ \cdots,\ \sigma(N_0)\}, \\
\sigma(1) < \sigma(2) < \cdots < \sigma(N_0).
\end{gathered}$$

Define a homeomorphism $\phi : T^N \to T^N$ by

$$(x_1, x_2, \cdots, x_i, \cdots, x_N) \mapsto (x_{\sigma(1)}, x_{\sigma(2)}, \cdots, x_{\sigma(i)}, \cdots, x_{\sigma(N)}).$$

Let

$$\beta(k) = \sum_{i=1}^{N_0} (-1)^{1-\tau_i(k)} \beta_{\sigma(i)},\ 1 \leqslant k \leqslant n = 2^{N_0}, \tag{4.1}$$

$$\gamma = \min_{1\leqslant k\leqslant n-1}(\beta(k+1)-\beta(k)), \tag{4.2}$$

$$\gamma_2 = \frac{1}{8\pi^2\lambda}\sum_{i=1}^{N}(\beta_i+M_0)^2, \tag{4.3}$$

$$T_1 = \sqrt{\frac{\gamma_1}{\gamma}},\ T_2 = \sqrt{\frac{\gamma}{\gamma_2}}. \tag{4.4}$$

Proposition 4.1. Assume that

$$\gamma > \sqrt{\gamma_1\gamma_2}. \tag{4.5}$$

Then

$$\gamma T^2 > \gamma_1 + \gamma_2 T^4,\ \forall T\in[T_1,T_2]. \tag{4.6}$$

Proof. By (4.4) and (4.5), we have $T_1 < T_2$. For each $T\in[T_1,T_2]$, we have

$$\gamma T^2 > \max\{\gamma_1,\ \gamma_2 T^4\}. \blacksquare$$

Proposition 4.2. *Let*

$$\Gamma = \left\{\frac{1}{2}\big(\beta(k+1)-\beta(k)\big)\ \middle|\ 1\leqslant k\leqslant n-1\right\}. \tag{4.7}$$

Then

$$\Gamma = \{\beta_{\sigma(N_0)}\}\bigcup\left\{\beta_{\sigma(i)} - \sum_{j=i+1}^{N_0}\beta_{\sigma(j)}\ \middle|\ 1\leqslant i\leqslant N_0-1\right\}.$$

Proof. Let

$$u(k) = (1-\tau_1(k),\cdots,1-\tau_{N_0}(k)).$$

Case 1. $N_0 = 1$. $n = 2^{N_0} = 2$. We have $k = 1$,

$$\begin{aligned} u(2) &= 0, \quad \beta(2) = \beta_1, \\ u(1) &= 1, \quad \beta(1) = -\beta_1. \end{aligned}$$

Then $\Gamma = \{\beta_1\}$.

Case 2. $N_0 \geqslant 2,\ n = 2^{N_0} \geqslant 4$.

$$\left.\begin{aligned} u(k+1) &= (u',0,\underbrace{1,\cdots,1}_{i}), \\ u(k) &= (u',1,\underbrace{0,\cdots,0}_{i}), \end{aligned}\right. \quad 0\leqslant i\leqslant N_0-1.$$

Then

$$\frac{1}{2}\big(\beta(k+1)-\beta(k)\big) = \begin{cases} \beta_{\sigma(N_0)}, & i = 0, \\ \beta_{\sigma(N_0-i)} - \displaystyle\sum_{j=N_0-i+1}^{N_0}\beta_{\sigma(j)}, & i = 1,\cdots,N_0-1. \blacksquare \end{cases}$$

4.2 Some estimates on potential energy part

Proposition 4.3. *For* $1 \leqslant k \leqslant 2^{N_0} - 1$, *we have*

$$V|_{\widetilde{M}_{k+1}} \leqslant \beta(k) < \beta(k+1) \leqslant V|_{\widetilde{R}^{-1}_{k+2}(\widetilde{S}_{k+1})}. \tag{4.8}$$

Proof. For $1 \leqslant s \leqslant m$, let

$$V_s(\xi_{m-s+1} \cdots, \xi_m) = \sum_{j=m-s+1}^{m} \beta_{\sigma(j)} \cos \xi_j, \tag{4.9}$$

and

$$h_s(k) = \sum_{j=m-s+1}^{m} (-1)^{1-\tau_j(k)} \beta_{\sigma(j)}. \tag{4.10}$$

Then

$$V_m|_{M_k^{(m)}} = V|_{\widetilde{M}_k}, \quad V_m\Big|_{R^{-1}_{m,k+1}\left(S_k^{(m)}\right)} = V\Big|_{\hat{R}^{-1}_{k+1}(\widetilde{S}_k)}, \quad h_m(k) = \beta(k)$$

and

$$h_m(k) - h_{m-1}(k) = (-1)^{1-\tau_1(k)} \beta_{\sigma(1)}. \tag{4.11}$$

By (3.18), we have

$$V_m\Big|_{R^{-1}_{m,k+1}\left(S_k^{(m)}\right)} \geqslant V_m\Big|_{O_k^{(m)}} \geqslant h_m(k).$$

We claim that

$$V_m|_{M_k^{(m)}} \leqslant h_m(k-1). \tag{4.12}$$

	$M_k^{(m)}$	$V_m\big|_{M_k^{(m)}} \leqslant$
$2 \leqslant k \leqslant 2^{m-1}$	$\{\pi\} \times M_k^{(m-1)}$	$-\beta_{\sigma(1)} + V_{m-1}\|_{M_k^{(m-1)}}$
$k = 2^{m-1} + 1$	$\{\pi\} \times T^{N-1}$	$-\beta_{\sigma(1)} + \sum_{j=2}^{m} \beta_{\sigma(j)} = \beta\left(2^{m-1}\right)$
$2^{m-1} + 2 \leqslant k \leqslant 2^m$	$\left(\{\pi\} \times T^{N-1}\right) \bigcup \left(S^1 \times M^{(m-1)}_{k-2^{m-1}}\right)$	$\beta_{\sigma(1)} + V_{m-1}\|_{M^{(m-1)}_{k-2^{m-1}}}$

Case 1. $2 \leqslant k \leqslant 2^{m-1}$. Assume that $V_{m-1}|_{M_k^{(m-1)}} \leqslant h_{m-1}(k-1)$. Then

$$V_m|_{M_k^{(m)}} \leqslant -\beta_{\sigma(1)} + V_{m-1}|_{M_k^{(m-1)}} \leqslant -\beta_{\sigma(1)} + h_{m-1}(k-1) = h_m(k-1).$$

Case 2. $k = 2^{m-1} + 1$. We have

$$\begin{gathered}
(k-1) - 1 = 2^{m-1} - 1 = 0 \cdot 2^{m-1} + 1 \cdot 2^{m-2} + \cdots + 1 \cdot 2^0, \\
\left(\tau_1(k-1), \tau_2(k-1), \cdots, \tau_m(k-1)\right) = (0, 1, 1, \cdots, 1), \\
h_{m-1}(k-1) = \sum_{j=2}^{m} (-1)^{1-\tau_j(k-1)} \beta_{\sigma(j)} = \sum_{j=2}^{m} \beta_{\sigma(j)}.
\end{gathered}$$

Then $V_m|_{M_k^{(m)}} \leqslant -\beta_{\sigma(1)} + \sum_{j=2}^{m} \beta_{\sigma(j)} = -\beta_{\sigma(1)} + h_{m-1}(k-1) = h_m(k-1)$.

Case 3. $2+2^{m-1} \leqslant k \leqslant 2^m$. We have

$$(k-1)-1 = 2^{m-1} + \left[\left(k-1-2^{m-1}\right)-1\right],\ 0 \leqslant \left(k-1-2^{m-1}\right)-1 \leqslant 2^{m-1}-2.$$

Then

$$\tau_j(k-1) = \begin{cases} 1, & j=1, \\ \tau_j\left(k-1-2^{m-1}\right), & 2 \leqslant j \leqslant m. \end{cases}$$

Assume that

$$V_{m-1}|_{M^{(m-1)}_{k-2^{m-1}}} \leqslant h_{m-1}\left(k-2^{m-1}-1\right).$$

We have

$$\begin{aligned} V_m\big|_{S^1\times M^{(m-1)}_{k-2^{m-1}}} &\leqslant \beta_{\sigma(1)} + V_{m-1}|_{M^{(m-1)}_{k-2^{m-1}}} \\ &\leqslant \beta_{\sigma(1)} + h_{m-1}\left(k-1-2^{m-1}\right) \\ &= \beta_{\sigma(1)} + h_{m-1}(k-1) = h_m(k-1). \end{aligned}$$

Note that

$$\begin{aligned} V_m\big|_{\{\pi\}\times T^{m-1}} &\leqslant -\beta_{\sigma(1)} + \sum_{j=2}^{m} \beta_{\sigma(j)} \leqslant \beta_{\sigma(1)} - \sum_{j=2}^{m} \beta_{\sigma(j)} \\ &\leqslant \beta_{\sigma(1)} + h_{m-1}(k-1) = h_m(k-1). \end{aligned}$$

Hence $V_m|_{M_k^{(m)}} \leqslant h_m(k-1)$. ∎

4.3 Proof of Theorem 2

For γ_1 define by (2.4) and $\beta(k)$ defined by (4.1), let

$$C_1(k) = T^{-1}\gamma_1 + T\beta(k) + f_v,\ 1 \leqslant k \leqslant n-1, \tag{4.13}$$

$$C_2(k) = T^{-1}2\pi^2|v|^2\lambda + T\beta(k+1) + f_v - T^3\gamma_2, \tag{4.14}$$

where

$$f_v = \int_0^T f(t)\cdot \frac{2\pi v t}{T}\mathrm{d}t.$$

By (4.6), we have

$$\begin{aligned} C_2(k) - C_1(k) &> T(\beta(k+1)-\beta(k)) - T^{-1}\gamma_1 - T^3\gamma_2 \\ &\geqslant T\cdot\gamma - T^{-1}\gamma_1 - T^3\gamma_2 > 0. \end{aligned} \tag{4.15}$$

Let

$$a_n = T^{-1}\gamma_1 + T\sum_{i=1}^{N_0} \beta_i + f_v + 1, \tag{4.16}$$

$$a_k = \frac{1}{2}(C_1(k)+C_2(k)),\ 1 \leqslant k \leqslant n-1. \tag{4.17}$$

By (4.15), we have

$$C_1(k) < a_k < C_2(k). \tag{4.18}$$

We claim that

$$a_1 < a_2 < \cdots < a_n. \tag{4.19}$$

and

$$\mathscr{L}|_{T^N\times\{0\}} < a_n,\ \mathscr{L}|_{\widetilde{M}_{k+1}\times\{0\}} < a_k < \mathscr{L}|_{p^{-1}\widetilde{R}^{-1}_{k+2}(\widetilde{S}_{k+1})},\ 1 \leqslant k \leqslant n-1, \tag{4.20}$$

Step 1. In fact, by (4.17) and (4.15), we have

$$\begin{aligned} a_{k+1} - a_k &= \frac{1}{2}(C_2(k+1) - C_2(k) + C_1(k+1) - C_1(k)) \\ &= \frac{T}{2}(\beta(k+2) - \beta(k+1) + \beta(k+1) - \beta(k)) \\ &\geqslant \frac{T}{2}(\gamma + \gamma) = T\gamma > 0. \end{aligned}$$

Step 2. We prove $\mathscr{L}|_{\widetilde{M}_{k+1}\times\{0\}} < a_k$. In fact, by the definition of $\mathscr{L}_2$, we have

$$\mathscr{L}_2|_{\widetilde{M}_{k+1}\times\{0\}} = TV|_{\widetilde{M}_{k+1}},\ \mathscr{L}_2|_{p^{-1}\widetilde{R}^{-1}_{k+2}(\widetilde{S}_{k+1})} = TV|_{\widetilde{R}^{-1}_{k+2}(\widetilde{S}_{k+1})}.$$

By Proposition 2.1 and Proposition 3.2, we have

$$\mathscr{L}_1|_{\widetilde{M}_{k+1}\times\{0\}} \leqslant T^{-1}\gamma_1,\ \mathscr{L}_2|_{\widetilde{M}_{k+1}\times\{0\}} \leqslant T\beta(k),\ \mathscr{L}_3|_{\widetilde{M}_{k+1}\times\{0\}} = f_v.$$

Then

$$\mathscr{L}|_{\widetilde{M}_{k+1}\times\{0\}} \leqslant T^{-1}\gamma_1 + T\beta(k) + f_v = C_1(k) < a_k.$$

Step 3. We prove $\mathscr{L}|_{p^{-1}\widetilde{R}^{-1}_{k+2}(\widetilde{S}_{k+1})} > a_k$. In fact, by Proposition 3.2, we have

$$\mathscr{L}_2|_{p^{-1}\widetilde{R}^{-1}_{k+2}(\widetilde{S}_{k+1})} \geqslant T\beta(k+1).$$

For each $x \in p^{-1}\widetilde{R}^{-1}_{k+2}(\widetilde{S}_{k+1})$, by Proposition 2.1, we have

$$\begin{aligned} \mathscr{L}(x) &\geqslant \sum_{i=1}^{N}\left[\frac{\lambda}{2}||\dot{x}_i||^2 - \frac{T\sqrt{T}}{2\pi}(\beta_i + M_0)||\dot{x}_i||\right] + T^{-1}2\pi^2|v|^2\lambda + \mathscr{L}_2(\bar{x}) + f_v \\ &\geqslant -\sum_{i=1}^{N}\frac{\left(\frac{T\sqrt{T}}{2\pi}(\beta_i + M_0)\right)^2}{2\lambda} + T^{-1}2\pi^2|v|^2\lambda + T\beta(k+1) + f_v \\ &= -T^3\gamma_2 + T^{-1}2\pi^2|v|^2\lambda + T\beta(k+1) + f_v = C_2(k) > a_k. \ \blacksquare \end{aligned}$$

References

[1] VI. Arnold, Mathematical Methods of Clasical Mechanics, New York, Springer-Verlag Inc. 1978. 75-93.

[2] J. Bellazzini, V. Benci, M. Ghimenti, Periodic orbits of a one-dimensional non-autonomous Hamiltonian system, *J. Differential Equations* **230** (2006) 275-294.

[3] A. Capozzi, D. Fortunato, A. Salvatore, Periodic solutions of Lagrangian systems with bounded potential, *J Math Anal Appl* **124** (1987) 482-494.

[4] K. C. Chang, Y. Long, E. Zehnder, Forced oscillation for the triple pendulum, *Analysis, et cetera*, Academic Press, Boston, 1990.

[5] K. C. Chang, *Infinite Dimensional Morse Theory and Multiple Solution Problems*, Birkhäuser, Boston, 1993.

[6] P. Felmer, Periodic solutions of spatially periodic Hamiltonian systems, *J. Diff. Eqns.* **98** (1992) 143–168.

[7] P. Felmer, Rotation type solutions for spatially periodic Hamiltonian systems, *Nonlinear Analysis, T.M.A.* **19** (1992) 409–425.

[8] G. Fournier, M. Willem, Multiple solutions of the forced double pendulum equation, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **6** (1989) 259–281.

[9] C. Golé, Monotone maps of $T^n \times \mathbb{R}^n$ and their periodic orbits, The geometry of Hamiltonian systems (ed. by T. Ratin), MSRI Publications 22 (Springer, New York) (1991) 341–366.

[10] F. Josellis, Ljusternik-Schnirelman theory for flows and periodic orbits for Hamiltonian systems on $T^n \times \mathbb{R}^n$, *Proc. London Math. Soc.* **68** (1994) 641–672.

[11] F. Josellis, Morse theory for forced oscillations of Hamiltonian systems on $T^n \times \mathbb{R}^n$, *J. Diff. Eqns.* **2** (1994) 360–384.

[12] R. Palais, The Lusternik-Schnirelman theory on Banach manifolds, *Topology* **5**(1966), 115-132.

[13] H. Qiao, Rotational solutions for Hamiltonian systems on $\mathbb{R}^n \times T^n$.

[14] P. Rabinowitz, On a class of functionals invariant under a action, *Trans. Am. math. Soc.* **310** (1988) 303–311.

[15] Paolo Roselli, A multiplicity result for the periodically forced N-pendulum with nonzero-mean valued forcings, *Nonlinear Anal. T.M.A.* **43** (2001) 1019-1041.

[16] E. Spanier, *Algebraic Topology*, McGraw-Hill, New York, 1966.

[17] G. Tarantello, Multiple forced oscillations for the N-pendulum equation, *Commun. Math. Phys.* **132** (1990) 499–517.